# **Diphenhydramine (Benadryl)**

## **Formulary**

**Douglas County Kansas EMS System** 

**November 2022** 

**Approved Provider:** AEMT, Paramedic

**Referenced Protocols:** Anaphylaxis and Allergic Reaction

Dosing and Administration	
ADULT Medications:	PEDIATRIC Medications: Refer to HandTevy
• 1.0-2.0 mg/kg IV, IO, IM (Max 50mg)	• 1.0-2.0 mg/kg IV, IO, IM (Max 50mg)

#### **Pharmacology and Actions:**

 An antihistamine which competitively blocks H-1 histamine receptors. Histamine can cause bronchoconstriction, vasodilatation, and capillary leak. It also has some anticholinergic (atropine-like) effects as well as direct CNS depressant effects

## **Indications:**

- Anaphylaxis
- Anaphylaxis with respiratory distress
- Anaphylactic shock
- Antidote for extrapyramidal reactions in phenothiazines and PCP

### **Contraindications:**

None

#### **Precautions:**

- In anaphylaxis, oxygen, epinephrine, fluids and Albuterol should be utilized prior to administration
- Due to anticholinergic effects, use with caution in patients with glaucoma, prostatism or peptic ulcer disease.

#### **Administration:**

Acceptable routes of administration include IV, IO and IM

### **Side Effects:**

- CNS depression
- Hypotension
- Thickening of bronchial secretions, tightness of chest and wheezing