Morphine Sulfate

Formulary

Douglas County Kansas EMS System

March 2021

Approved Provider: Paramedic

Referenced Protocols: Pain/Anxiety Management

Dosing and Administration	
ADULT Medications:	PEDIATRIC Medications: Refer to HandTevy
 Analgesia: 2-4 mg IV/IO/IM, q 5 min (max 10 mg) 	• Analgesia: 0.1 mg/kg IV/IO/IM, q 5 min (max 10 mg)

Pharmacology and Actions:

- Acts as an opiate agonist
- Alters the patient's perception of pain and acts as an analgesic by stimulating opiate receptor sites
- Produces venous vasodilatation

Indications:

- For the treatment of pain and anxiety associated with AMI
- Useful to relieve pain associated with burns and isolated trauma
- As an adjunct in the treatment of pulmonary edema

Contraindications:

- Respiratory distress or compromise (Asthma or COPD)
- Altered mental status
- Sensitivity/allergy to Morphine
- Hypotension and suspected shock

Precautions:

- This drug causes decreased respiratory drive, and as such should only be given while monitoring pulse oximetry and closely monitoring respiratory status
- Pregnancy
- Drug/alcohol intoxication
- Age less than 12 months
- Stimulates Vomiting

Administration:

- For specific doses, refer to applicable protocol
- May be administered IV only

Side Effects:

- CNS- Depression, Convulsions in children
- CV- Bradycardia, Decrease Blood pressure
- Respiratory- Depression, Wheezing
- G/I- Nausea/Vomiting